

Gravity Induced New Topological Phase in Optics

Partha Ghose and M. K. Samal

S. N. Bose National Centre for Basic Sciences, Block JD, Sector III, Salt Lake, Kolkata 700 098

It is shown that both classical and quantum light can acquire a topological phase shift induced by classical gravity, and the latter is detectable in a laboratory-scale experiment.

Ever since the Aharonov-Bohm (AB) effect [1] was theoretically predicted, there has been the assumption that it is a purely quantum mechanical effect without any classical counterpart. The electrodynamic potentials themselves suddenly acquired a physical significance that was lacking in classical electrodynamics. However, Berry [2] has shown that the Pancharatnam phase [3] for polarised classical light is analogous to the AB phase. It is the purpose of this paper to show that classical light can also exhibit a topological phase similar to the AB phase and which is gravity-induced. This type of effect holds in classical physics as long as (a) the classical system under consideration has a wave-like character (as in electrodynamics) and (b) the space is not simply connected. A very important consequence of this classical optical analogue of the AB effect implies that the classical gravitational ‘potential’ has a physical significance analogous to that of the electromagnetic potentials in quantum mechanics. In fact, all potentials will acquire physical significance in spaces that are not simply connected.

To see the classical optical analogue of the scalar AB effect let us consider a typical interference experiment with a Mach-Zehnder interferometer, a pulsed source of classical light (with Poissonian statistics) of energy E (and therefore of effective mass E/c^2) and a thin spherical shell of mass M and radius R placed in the path of one of the two beams inside the interferometer. The spherical shell has two small holes diametrically opposite each other that allow the light pulses to pass through its field-free interior. The interaction Hamiltonian H_{int} inside the shell can be written as

$$H_{int} = V_g \frac{E}{c^2} = -\frac{GM}{R} \frac{E}{c^2} = -\frac{GM}{R} \left(\frac{4\pi R^3}{3c^2} \frac{\epsilon_0 \mathbf{E} \cdot \mathbf{E}}{2} \right) \equiv -\frac{4}{3} \pi R^3 \cdot \frac{1}{2} \epsilon_g \mathbf{E} \cdot \mathbf{E} \quad (1)$$

where ϵ_g is the effective gravity-induced permittivity seen by the light with electric vector $\mathbf{E}$ although it does not experience any gravitational force. It follows from this that the effective refractive index is given by

$$n = \sqrt{\epsilon_0 + \epsilon_g} \approx \sqrt{\epsilon_0} \left(1 + \frac{GM}{2Rc^2} \right) \quad (2)$$

since $\epsilon_g \ll \epsilon_0$. Therefore light emerging from the shell will experience a gravity-induced topological phase shift given by

$$\phi_{cl} = \frac{2\pi}{\lambda} (2R)(n - \sqrt{\epsilon_0}) \approx \frac{2\pi GM}{\lambda c^2} \sqrt{\epsilon_0} \quad (3)$$

where λ is the wavelength of light.

The constant potential $V = GM/R$ cannot be gauged away because the spaces inside and outside the sphere are not simply connected. Therefore, this phase shift will be observable, and is the new optical analogue of the scalar AB shift pointed out by Aharonov and Bohm in their original paper.

The quantum optical analogue of the scalar AB effect can be studied by writing the classical Maxwell theory of electrodynamics in the Schrödinger form [5]

$$i\hbar \frac{\partial(\gamma\psi)}{\partial t} = (H_0 + H_{int})(\gamma\psi) \quad (4)$$

with $\psi^T = \frac{1}{\sqrt{m}}(-E_x, -E_y, -E_z, H_x, H_y, H_z, -mA_x, -mA_y, -mA_z, mA_0)$ and

$$H_0 = -i\hbar \tilde{\beta}_i \partial_i \psi \quad (5)$$

where $\tilde{\beta}_i = \beta_0 \beta_i - \beta_i \beta_0$, the β 's being the (10×10) Kemmer-Duffin-Petiau matrices satisfying the algebra

$$\beta_\mu \beta_\nu \beta_\lambda + \beta_\lambda \beta_\nu \beta_\mu = \beta_\mu g_{\nu\lambda} + \beta_\lambda g_{\nu\mu} \quad (6)$$

and

$$\gamma \beta_\mu + \beta_\mu \gamma = \beta_\mu \quad (7)$$

The matrix γ is diagonal with the first six diagonal elements unity and the last four diagonal elements zero. Equation (4) must be supplemented by the gauge-invariant first-class constraint equation

$$i\hbar\beta_i\beta_0^2\partial_i\psi + m(1 - \beta_0^2)\gamma\psi = 0 \quad (8)$$

Equations (4) and (8) together are equivalent to the Lorentz invariant equation

$$i\hbar\beta_\mu\partial^\mu\psi + m\gamma\psi = 0 \quad (9)$$

which implies the massless second-order equation

$$\square(\gamma\psi) = 0 \quad (10)$$

The Schrödinger equation (4) translates into the Maxwell equations

$$\begin{aligned} \text{curl } \mathbf{H} &= \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t} \\ \text{curl } \mathbf{E} &= -\frac{1}{c} \frac{\partial \mathbf{H}}{\partial t} \end{aligned} \quad (11)$$

and the constraint equation (8) into the equations

$$\begin{aligned} \text{div } \mathbf{E} &= 0 \\ \mathbf{H} &= \text{curl } \mathbf{A} \end{aligned} \quad (12)$$

This set of four equations together constitute the complete set of Maxwell equations. This shows that although the quantum mechanical equation (4) involves $\hbar$, the Maxwell equations which are contained in it do not. Hence it is not possible to say whether the Maxwell equations by themselves describe a classical or quantum system. It is nowadays well known that the quantum nature of radiation is difficult to observe, and depends crucially on how light is produced. In most production processes the light produced has Poissonian statistics and is indistinguishable from classical light. It is only when special care is taken to produce Fock states of light (as in parametric down-conversion of laser light) or squeezed states that the quantum nature of radiation can be observed. *Hence the production process constitutes the additional constraint that renders the Maxwell equations either classical or quantum.*

There are three features of this formalism that must be pointed out. First, there is a conserved four-vector current s_μ in this theory whose time component s_0 is positive definite, and can be identified with the probability current [6]. It is associated with the energy flux and energy density. This makes a consistent quantum mechanical interpretation of this formalism possible. Second, the formalism describes a massless wave/particle in spite of the mass parameter m which is essential for dimensional reasons. Nevertheless, the theory is gauge invariant under the local transformations $\psi \rightarrow \psi + (1 - \gamma)\chi$ where χ is an arbitrary function of space and time. In fact, this mass parameter can be expressed in terms of the fundamental constants of the theory by the relation $m = s\hbar\omega/c^2$ where ω is the frequency of the photon and $s = 2$ is the spin degeneracy. This aspect of the formalism has not been pointed out before. Third, since the classical Maxwell equations (11) and (12) are contained in this quantum mechanical formalism but do not involve any dimensional mass parameter, it is clear that *the classical angular momentum as well as the classical polarization states are totally independent of the quantum mechanical spin σ .* We are not aware of any other formalism in which this distinction between the quantum mechanical spin of the photon and the angular momentum of the classical electromagnetic field is so clearly expressed.

If one considers the same type of experiment discussed in the previous section but with a pulse of laser light of energy $E \approx \bar{N}\hbar\nu$ where $\bar{N}$ is the average number of photons in the pulse around which the distribution is peaked, then the phase shift predicted by the Schrödinger equation (4) is

$$\phi_{qm} = \frac{H_{int}t}{\hbar} = \frac{4\pi G M \bar{N}}{\lambda c^2} \quad (13)$$

since $\epsilon_0 = 1$. The advantage of using a laser pulse is that the phase shift is enhanced by the factor $\bar{N}$ compared to a single photon state. Using a typical pico second pulsed laser with wavelength $\approx 5000\text{\AA}$ one gets $\bar{N} \approx 10^7$. This enhancement is not possible in the classical limit which corresponds to $\bar{N} \ll 1$. This can also be seen from the fact that an increase in the intensity of classical light does not enhance the refractive index (hence the phase shift) because of the linearity of the vacuum.

The question is whether the predicted phase shifts can be detected. For classical light of wavelength $\approx 5000\text{\AA}$ one needs a thin spherical shell of mass $\approx 10^{18}$ kg to produce a phase shift of the order of a milliradian. This kind of mass can only be realized in astrophysical conditions. Nevertheless a laboratory-scale experiment could still be possible by compensating for the large mass through an increase of the effective optical path (through the sphere) by repeatedly circulating the same light pulse as shown in the figure. This is equivalent to introducing a large winding number n_w .




FIG. 1. A modified Mach-Zehnder set-up

The figure shows a modified Mach-Zehnder set-up having additional lossless mirrors m_{ij} with m_{12} , m_{13} , m_{22} , m_{23} fixed and m_{11} , m_{14} , m_{21} , m_{24} movable. m_{11} and m_{21} are inserted from the beginning and removed after n cycles are completed by the light pulses. m_{14} and m_{24} are inserted after the light pulses enter the sphere and in time for them to circulate as shown.

Note that the light pulses acquire a dynamical phase while circulating in the gravitational field between the mirrors. If the paths between the fixed mirrors m_{12} , m_{22} and m_{13} , m_{23} are kept very close, the dynamical phases picked up along these paths will be nearly the same, and hence will cancel one another in the interference pattern. The only additional dynamical phase in the lower circuit will come from the path between the mirrors m_{24} and m_{21} . This can be made significantly small compared to the topological phase acquired in the upper circuit by arranging the distance between the mirror $M1$ and the beam-splitter $BS1$ large enough compared to the radius R of the spherical shell.

A simple calculation shows that for a thin spherical shell of radius ≈ 3.3 m, thickness 10 cm and mass $\approx 10^5$ kg one needs a winding number $n_w = 10^{12}$ which would require a duration of the order of 15 hours. The problem in actually performing the suggested experiment would be maintaining the required stability of the entire set-up as well as the lossless character of the mirrors to detect the phase-shift of the order of a milliradian over such a long duration.

As we have seen, unlike the classical situation, the use of a pulsed laser has the advantage of enhancing the phase shift by a factor of $\bar{N} \approx 10^7$. This enables a reduction in the dimensions of the spherical shell to radius ≈ 1.5 m, thickness 1 cm and mass $\approx 3 \times 10^3$ kg for a winding number $n_w = 10^6$. This can be achieved in a time interval of the order of 0.1 sec which makes the experiment feasible.

The new result of this work, is that both *classical* and quantum light can acquire a topological phase that is induced by *classical* gravity and the latter can be measured in a laboratory-scale experiment. One important implication of this is that the classical gravitational potential acquires a new physical significance in spaces that are not simply connected.

We thank DST, Govt. of India for financial support to carry out this work and A. Greentree for helpful correspondence.

-
- [1] Y. Aharonov and D. Bohm, *Phys. Rev.* **115**, 485, 1059.
 - [2] M. V. Berry, *Jour. Mod. Optics* **34**, 1401, 1987.
 - [3] S. Pancharatnam, *Proc. Ind. Acad. of Sciences* **XLIV** 247, 1956.
 - [4] P. Ghose, quant-ph/0104104.
 - [5] Harish-Chandra, *Proc. Roy. Soc. A* **186**, 502, 1946; N. Kemmer, *Proc. Roy. Soc. A* **173**, 91, 1939.
 - [6] P. Ghose, *Found. of Phys.* **26**, 1441, 1996; P. Ghose, D. Home and M. N. Sinha Roy, *Phys. Lett. A* **183**, 267, 1993.